

# First-order logic Resolution

Mario Alviano

University of Calabria, Italy

A.Y. 2018/2019

- 1 First-order resolution
  - Unification
  - From propositional to first-order resolution
  
- 2 Exercises

How to generalize propositional resolution to first-order?

Biggest obstacle: When two atoms are “equal”?

$$\forall x (H(x) \rightarrow M(x)) \wedge H(\text{socrates})$$

- Resolution works on Skolem CNF formulas!
  - 1  $\{\neg H(x), M(x)\}$
  - 2  $\{H(\text{socrates})\}$
- Can we derive  $M(\text{socrates})$ ?
  - $H(x)$  is different from  $H(\text{socrates})$
- Can we substitute  $x$  by  $\text{socrates}$ ?
- Let's formalize this idea!

# Composition of substitutions

Recall that  $[t_1/x_1, \dots, t_n/x_n]$  denotes the (simultaneous) substitution of  $x_1, \dots, x_n$  with  $t_1, \dots, t_n$ .

## Definition

Let  $\sigma = [t_1/x_1, \dots, t_n/x_n]$ ,  $\vartheta = [s_1/y_1, \dots, s_m/y_m]$  be substitutions. The composition of  $\sigma$  and  $\vartheta$ , denoted  $\sigma \circ \vartheta$  (or  $\sigma\vartheta$ ), is obtained from  $[t_1\vartheta/x_1, \dots, t_n\vartheta/x_n, s_1/y_1, \dots, s_m/y_m]$

- by removing each  $t_i\vartheta/x_i$  such that  $t_i\vartheta = x_i$ , and
- by removing each  $s_j/y_j$  such that  $y_j \in \{x_1, \dots, x_n\}$ .

Just remember that substitutions are functions!

## Properties

- $\sigma[] = []\sigma = \sigma$
- $(\sigma\vartheta)\rho = \sigma(\vartheta\rho)$
- $\sigma\vartheta \neq \vartheta\sigma$
- $(t\sigma)\vartheta = t(\sigma\vartheta) = t\sigma\vartheta$
- $(\phi\sigma)\vartheta = \phi(\sigma\vartheta) = \phi\sigma\vartheta$
- $(E\sigma)\vartheta = E(\sigma\vartheta) = E\sigma\vartheta$

## Definition

Let  $E_1, E_2$  be first-order expressions (i.e., terms or formulas).

- A substitution  $\sigma$  is a **unifier** of  $E_1, E_2$  if  $E_1\sigma = E_2\sigma$
- A unifier  $\sigma$  is a **most general unifier (mgu)** if for any unifier  $\vartheta$  of  $E_1, E_2$  it holds that  $\vartheta = \sigma\rho$  for some substitution  $\rho$

## Properties

- If  $E_1, E_2$  are unifiable then an mgu exists
- Moreover, the mgu is unique modulo variable renaming

Let  $\perp$  denote failure.

- $unify(P(t_1, \dots, t_n), Q(s_1, \dots, s_m)) = \perp$  if  $P \neq Q$  or  $n \neq m$
- $unify(P(t_1, \dots, t_n), P(s_1, \dots, s_n)) = unify((t_1, \dots, t_n), (s_1, \dots, s_n))$
- $unify(), () = []$
- $unify((t_1, \dots, t_n), (s_1, \dots, s_n)) = \perp$  if  $unify(t_1, s_1) = \perp$
- $unify((t_1, \dots, t_n), (s_1, \dots, s_n)) = \sigma \circ unify((t_2\sigma, \dots, t_n\sigma), (s_2\sigma, \dots, s_n\sigma))$  where  $\sigma = unify(t_1, s_1)$  and  $\sigma \neq \perp$
- $unify(f(t_1, \dots, t_n), g(s_1, \dots, s_m)) = \perp$  if  $f \neq g$  or  $n \neq m$
- $unify(f(t_1, \dots, t_n), f(s_1, \dots, s_n)) = unify((t_1, \dots, t_n), (s_1, \dots, s_n))$
- $unify(t, x) = unify(x, t)$  if  $t$  is not a variable
- $unify(x, t) = [t/x]$  if  $x \notin free(t)$
- $unify(x, t) = \perp$  if  $x \in free(t)$  and  $x \neq t$
- $unify(x, x) = []$

## Examples

- 1  $unify(P(x, f(x)), P(y, f(g(b))))$
- 2  $unify(A(x, y), A(y, f(z)))$
- 3  $unify(B(a, y, f(y)), B(z, z, u))$
- 4  $unify(A(x, g(x)), A(y, y))$

# Resolvent and factorization

## Resolvent

Let  $C_1, C_2$  be two clauses.

- Assume two variable renaming substitutions  $\sigma_1$  and  $\sigma_2$  such that  $C_1\sigma_1$  and  $C_2\sigma_2$  do not share variables.
- If  $A_1 \in C_1\sigma_1$  and  $\neg A_2 \in C_2\sigma_2$  such that  $A_1$  and  $A_2$  are unifiable with mgu  $\vartheta$ , then

$$((C_1\sigma_1 \setminus \{A_1\}) \cup (C_2\sigma_2 \setminus \{\neg A_2\}))\vartheta$$

is a **resolvent** of  $C_1$  and  $C_2$ .

## Factorization

Given a clause  $C$ , if there are literals  $A_1, A_2$  (or  $\neg A_1, \neg A_2$ ) of  $C$  such that  $A_1$  and  $A_2$  are unifiable with mgu  $\vartheta$ , then  $C\vartheta$  is a **factor** of  $C$ .




## Derivation

Given a set  $\Gamma$  of clauses, a derivation by resolution of a clause  $C$  from  $\Gamma$ , denoted  $\Gamma \vdash_R C$ , is a sequence  $C_1, \dots, C_n$  such that  $C_n = C$  and for each  $C_i$  ( $1 \leq i \leq n$ ) we have

- 1  $C_i \in \Gamma$ , or
- 2  $C_i$  is a resolvent of  $C_j$  and  $C_k$ , where  $j < i$  and  $k < i$ , or
- 3  $C_i$  is a factor of  $C_j$ , where  $j < i$ .

We can now use all other definitions and claims from propositional resolution!



- 1 Find the most general unifier for  $A(r(c, x), r(z, z))$  and  $A(y, y)$  (if it exists)
- 2 Decide by means of resolution whether the following formulas (in SCNF) are satisfiable:
  - 1  $\{R(x, g(y))\}, \{\neg R(v, v), S(v)\}, \{\neg S(g(c))\}$
  - 2  $\{R(x, g(x))\}, \{\neg R(z, z), S(z)\}, \{\neg S(g(x))\}$
  - 3  $\{R(x, y), R(z, c), S(z)\}, \{\neg R(f(u), u), \neg R(v, w)\}, \{\neg S(f(d))\}$

1 Use resolution to prove that

- $\forall x \forall y \forall z (P(x, y) \wedge P(y, z) \rightarrow P(x, z))$
- $\forall x \forall y (P(x, y) \rightarrow P(y, x))$

entail

- $\forall x \forall y \forall z (P(x, y) \wedge P(z, y) \rightarrow P(x, z))$

2 Use resolution to prove that

- $\forall x A(a, x, x)$
- $\forall x \forall y \forall z (A(x, y, z) \rightarrow A(x, s(y), s(z)))$
- $\forall x \forall y \forall z (A(x, y, z) \rightarrow A(y, x, z))$

entail

- $\exists x A(s(s(a)), s(s(s(a))), x)$

Exercises 6.18 and 6.19 from *Logica a Informatica*.

1 Prove using resolution that 1 entails 2:

1 All students are citizens

2 Votes by students are votes by citizens

**Hint:** Use  $Student(x)$ ,  $Citizen(x)$  and  $Votes(x, y)$  for “x is a student”, “x is a citizen” and “x votes y”, respectively.

2 Check whether 3 is a logical consequence of 1 and 2:

1 Every lion chases some gazelle

2 Every gazelle fears anyone chasing it


3 Every gazelle fears someone

In case your answer is negative, provide a Herbrand model.

Exercise 6.20 from *Logica a Informatica*.

- 1 Check whether 4 is a logical consequence of the other sentences:
  - 1 Policemen searched all those who have stepped off the plane and were not member of the crew
  - 2 Some thieves (one or more) stepped off the plane and were searched only by thieves
  - 3 No thief was a member of the crew
  - 4 Some policemen (one or more) were thieves

In case your answer is negative, provide a Herbrand model.



END OF THE  
LECTURE