

First-order logic

Normal forms and Herbrand theory

Mario Alviano

University of Calabria, Italy

A.Y. 2018/2019

- 1 Normal forms
 - Prenex Normal Form
 - Skolemization
- 2 Herbrand theory
 - Intuition
 - Main statement
- 3 Exercises

- 1 Normal forms
 - Prenex Normal Form
 - Skolemization
- 2 Herbrand theory
 - Intuition
 - Main statement
- 3 Exercises

- 1 Normal forms
 - Prenex Normal Form
 - Skolemization
- 2 Herbrand theory
 - Intuition
 - Main statement
- 3 Exercises

Prenex Normal Form

Formulas of the following type are in Prenex Normal Form:

$$Q_1 x_1 \cdots Q_n x_n \phi$$

where

- $Q_i \in \{\forall, \exists\}$ for $i = 1, \dots, n$
 - $Q_1 x_1 \cdots Q_n x_n$ is called **quantifier prefix**
- ϕ is a quantifier-free formula
 - ϕ is called **matrix**

Prenex Normal Form

Formulas of the following type are in Prenex Normal Form:

$$Q_1 x_1 \cdots Q_n x_n \phi$$

where

- $Q_i \in \{\forall, \exists\}$ for $i = 1, \dots, n$
 - $Q_1 x_1 \cdots Q_n x_n$ is called **quantifier prefix**
- ϕ is a quantifier-free formula
 - ϕ is called **matrix**

Theorem

For each wff ϕ there is an equivalent ϕ^{PNF} in prenex normal form.

Prenex Normal Form

Formulas of the following type are in Prenex Normal Form:

$$Q_1 x_1 \cdots Q_n x_n \phi$$

where

- $Q_i \in \{\forall, \exists\}$ for $i = 1, \dots, n$
 - $Q_1 x_1 \cdots Q_n x_n$ is called **quantifier prefix**
- ϕ is a quantifier-free formula
 - ϕ is called **matrix**

Theorem

For each wff ϕ there is an equivalent ϕ^{PNF} in prenex normal form.

Intuitively, move quantifiers outside, i.e., up on the tree structure

Negation Normal Form

- 1 Compute the prenex normal form
- 2 Compute the negation normal form of the matrix

Negation, Conjunctive and Disjunctive Normal Form

Negation Normal Form

- 1 Compute the prenex normal form
- 2 Compute the negation normal form of the matrix

Conjunctive Normal Form

- 1 Compute the prenex normal form
- 2 Compute the conjunctive normal form of the matrix

Negation, Conjunctive and Disjunctive Normal Form

Negation Normal Form

- 1 Compute the prenex normal form
- 2 Compute the negation normal form of the matrix

Conjunctive Normal Form

- 1 Compute the prenex normal form
- 2 Compute the conjunctive normal form of the matrix

Disjunctive Normal Form

- 1 Compute the prenex normal form
- 2 Compute the disjunctive normal form of the matrix

- 1 Normal forms
 - Prenex Normal Form
 - Skolemization
- 2 Herbrand theory
 - Intuition
 - Main statement
- 3 Exercises

- Can we further simplify the structure of a wff?

- Can we further simplify the structure of a wff?
- Can we remove existential quantifiers?

- Can we further simplify the structure of a wff?
- Can we remove existential quantifiers?
- Yes, by introducing **Skolem functions**!

- Can we further simplify the structure of a wff?
- Can we remove existential quantifiers?
- Yes, by introducing **Skolem functions**!
- However, equivalence will not hold anymore

- Can we further simplify the structure of a wff?
- Can we remove existential quantifiers?
- Yes, by introducing **Skolem functions**!
- However, equivalence will not hold anymore
- Still we will preserve satisfiability!

- Can we further simplify the structure of a wff?
- Can we remove existential quantifiers?
- Yes, by introducing **Skolem functions**!
- However, equivalence will not hold anymore
- Still we will preserve satisfiability!

Intuition

Consider $\exists x A(x)$

- Can we further simplify the structure of a wff?
- Can we remove existential quantifiers?
- Yes, by introducing **Skolem functions**!
- However, equivalence will not hold anymore
- Still we will preserve satisfiability!

Intuition

Consider $\exists x A(x)$

- There is x such that $A(x)$ is true

- Can we further simplify the structure of a wff?
- Can we remove existential quantifiers?
- Yes, by introducing **Skolem functions**!
- However, equivalence will not hold anymore
- Still we will preserve satisfiability!

Intuition

Consider $\exists x A(x)$

- There is x such that $A(x)$ is true
- Replace x with a **fresh** constant c

- Can we further simplify the structure of a wff?
- Can we remove existential quantifiers?
- Yes, by introducing **Skolem functions**!
- However, equivalence will not hold anymore
- Still we will preserve satisfiability!

Intuition

Consider $\exists x A(x)$

- There is x such that $A(x)$ is true
- Replace x with a **fresh** constant c
- The Skolem formula will be $A(c)$

- Can we further simplify the structure of a wff?
- Can we remove existential quantifiers?
- Yes, by introducing **Skolem functions**!
- However, equivalence will not hold anymore
- Still we will preserve satisfiability!

Intuition

Consider $\exists x A(x)$

- There is x such that $A(x)$ is true
- Replace x with a **fresh** constant c
- The Skolem formula will be $A(c)$

Consider $\forall x \exists y B(x, y)$

- Can we further simplify the structure of a wff?
- Can we remove existential quantifiers?
- Yes, by introducing **Skolem functions**!
- However, equivalence will not hold anymore
- Still we will preserve satisfiability!

Intuition

Consider $\exists x A(x)$

- There is x such that $A(x)$ is true
- Replace x with a **fresh** constant c
- The Skolem formula will be $A(c)$

Consider $\forall x \exists y B(x, y)$

- Can we replace y by a constant d ?

- Can we further simplify the structure of a wff?
- Can we remove existential quantifiers?
- Yes, by introducing **Skolem functions**!
- However, equivalence will not hold anymore
- Still we will preserve satisfiability!

Intuition

Consider $\exists x A(x)$

- There is x such that $A(x)$ is true
- Replace x with a **fresh** constant c
- The Skolem formula will be $A(c)$

Consider $\forall x \exists y B(x, y)$

- Can we replace y by a constant d ?
 - \mathcal{A} such that $D_{\mathcal{A}} = \mathbb{N}^+$ and $B^{\mathcal{A}} = \{(x, y) \mid x \text{ divides } y\}$

- Can we further simplify the structure of a wff?
- Can we remove existential quantifiers?
- Yes, by introducing **Skolem functions**!
- However, equivalence will not hold anymore
- Still we will preserve satisfiability!

Intuition

Consider $\exists x A(x)$

- There is x such that $A(x)$ is true
- Replace x with a **fresh** constant c
- The Skolem formula will be $A(c)$

Consider $\forall x \exists y B(x, y)$

- Can we replace y by a constant d ?
 - \mathcal{A} such that $D_{\mathcal{A}} = \mathbb{N}^+$ and $B^{\mathcal{A}} = \{(x, y) \mid x \text{ divides } y\}$
 - Is d a multiple of every natural number?

- Can we further simplify the structure of a wff?
- Can we remove existential quantifiers?
- Yes, by introducing **Skolem functions**!
- However, equivalence will not hold anymore
- Still we will preserve satisfiability!

Intuition

Consider $\exists x A(x)$

- There is x such that $A(x)$ is true
- Replace x with a **fresh** constant c
- The Skolem formula will be $A(c)$

Consider $\forall x \exists y B(x, y)$

- Can we replace y by a constant d ?
 - \mathcal{A} such that $D_{\mathcal{A}} = \mathbb{N}^+$ and $B^{\mathcal{A}} = \{(x, y) \mid x \text{ divides } y\}$
 - Is d a multiple of every natural number?
- Let's replace y by a function $f(x)$

Skolem Normal Form

Let ϕ be a wff. ϕ^{SNF} is obtained as follows:

- 1 Compute the prenex normal form of ϕ

$$\gamma := Q_1 x_1 \cdots Q_n x_n \psi$$

- 2 If γ has no existential quantifier then **stop**
- 3 Let $i \in \{1, \dots, n\}$ be the first existential quantifier
- 4 Modify γ :
 - Remove the quantifier $Q_i x_i$
 - Replace x_i by $f(x_1, \dots, x_{i-1})$
where f is a fresh function symbol of arity $i - 1$
- 5 Go to **2**

Skolem Normal Form

Let ϕ be a wff. ϕ^{SNF} is obtained as follows:

- 1 Compute the prenex normal form of ϕ

$$\gamma := Q_1 x_1 \cdots Q_n x_n \psi$$

- 2 If γ has no existential quantifier then **stop**
- 3 Let $i \in \{1, \dots, n\}$ be the first existential quantifier
- 4 Modify γ :
 - Remove the quantifier $Q_i x_i$
 - Replace x_i by $f(x_1, \dots, x_{i-1})$
where f is a fresh function symbol of arity $i - 1$
- 5 Go to **2**

Functions introduced by the Skolemization are called
Skolem functions

Theorem

Let ϕ be a wff.

ϕ is satisfiable if and only if ϕ^{SNF} is satisfiable.

Theorem

Let ϕ be a wff.

ϕ is satisfiable if and only if ϕ^{SNF} is satisfiable.

Corollary

Let ϕ be a wff.

ϕ is satisfiable if and only if $Ex(\phi)^{SNF}$ is satisfiable.

Theorem

Let ϕ be a wff.

ϕ is satisfiable if and only if ϕ^{SNF} is satisfiable.

Corollary

Let ϕ be a wff.

ϕ is satisfiable if and only if $Ex(\phi)^{SNF}$ is satisfiable.

Let's consider only closed formulas!

- 1 Normal forms
 - Prenex Normal Form
 - Skolemization
- 2 Herbrand theory
 - Intuition
 - Main statement
- 3 Exercises

- 1 Normal forms
 - Prenex Normal Form
 - Skolemization
- 2 Herbrand theory
 - Intuition
 - Main statement
- 3 Exercises

Herbrand intuition

So many models! Even for a simple formula like $P(c)$ there are infinitely many structures and models

Herbrand intuition

So many models! Even for a simple formula like $P(c)$ there are infinitely many structures and models

$$\mathcal{A}_1 \not\models P(c)$$

$$D_{\mathcal{A}_1} = \{a\}$$

$$c^{\mathcal{A}_1} = a$$

$$P^{\mathcal{A}_1}(a) = 0$$

Herbrand intuition

So many models! Even for a simple formula like $P(c)$ there are infinitely many structures and models

$\mathcal{A}_1 \not\models P(c)$

$D_{\mathcal{A}_1} = \{a\}$

$c^{\mathcal{A}_1} = a$

$P^{\mathcal{A}_1}(a) = 0$

$\mathcal{A}_2 \models P(c)$

$D_{\mathcal{A}_2} = \{a\}$

$c^{\mathcal{A}_2} = a$

$P^{\mathcal{A}_2}(a) = 1$

Herbrand intuition

So many models! Even for a simple formula like $P(c)$ there are infinitely many structures and models

$$\mathcal{A}_1 \not\models P(c)$$

$$\begin{aligned} D_{\mathcal{A}_1} &= \{a\} \\ c^{\mathcal{A}_1} &= a \\ P^{\mathcal{A}_1}(a) &= 0 \end{aligned}$$

$$\mathcal{A}_2 \models P(c)$$

$$\begin{aligned} D_{\mathcal{A}_2} &= \{a\} \\ c^{\mathcal{A}_2} &= a \\ P^{\mathcal{A}_2}(a) &= 1 \end{aligned}$$

$$\mathcal{A}_3 \not\models P(c)$$

$$\begin{aligned} D_{\mathcal{A}_3} &= \{b\} \\ c^{\mathcal{A}_3} &= b \\ P^{\mathcal{A}_3}(b) &= 0 \end{aligned}$$

Herbrand intuition

So many models! Even for a simple formula like $P(c)$ there are infinitely many structures and models

$\mathcal{A}_1 \not\models P(c)$

$D_{\mathcal{A}_1} = \{a\}$
 $c^{\mathcal{A}_1} = a$
 $P^{\mathcal{A}_1}(a) = 0$

$\mathcal{A}_2 \models P(c)$

$D_{\mathcal{A}_2} = \{a\}$
 $c^{\mathcal{A}_2} = a$
 $P^{\mathcal{A}_2}(a) = 1$

$\mathcal{A}_3 \not\models P(c)$

$D_{\mathcal{A}_3} = \{b\}$
 $c^{\mathcal{A}_3} = b$
 $P^{\mathcal{A}_3}(b) = 0$

$\mathcal{A}_4 \models P(c)$

$D_{\mathcal{A}_4} = \{b\}$
 $c^{\mathcal{A}_4} = b$
 $P^{\mathcal{A}_4}(b) = 1$

Herbrand intuition

So many models! Even for a simple formula like $P(c)$ there are infinitely many structures and models

$\mathcal{A}_1 \not\models P(c)$

$D_{\mathcal{A}_1} = \{a\}$
 $c^{\mathcal{A}_1} = a$
 $P^{\mathcal{A}_1}(a) = 0$

$\mathcal{A}_2 \models P(c)$

$D_{\mathcal{A}_2} = \{a\}$
 $c^{\mathcal{A}_2} = a$
 $P^{\mathcal{A}_2}(a) = 1$

$\mathcal{A}_3 \not\models P(c)$

$D_{\mathcal{A}_3} = \{b\}$
 $c^{\mathcal{A}_3} = b$
 $P^{\mathcal{A}_3}(b) = 0$

$\mathcal{A}_4 \models P(c)$

$D_{\mathcal{A}_4} = \{b\}$
 $c^{\mathcal{A}_4} = b$
 $P^{\mathcal{A}_4}(b) = 1$

$\mathcal{A}_5 \not\models P(c)$

$D_{\mathcal{A}_5} = \{a, b\}$
 $c^{\mathcal{A}_5} = b$
 $P^{\mathcal{A}_5}(b) = 0$
 $P^{\mathcal{A}_5}(a) = 0$
(or $P^{\mathcal{A}_5}(a) = 1$)

Herbrand intuition

So many models! Even for a simple formula like $P(c)$ there are infinitely many structures and models

$\mathcal{A}_1 \not\models P(c)$

$D_{\mathcal{A}_1} = \{a\}$
 $c^{\mathcal{A}_1} = a$
 $P^{\mathcal{A}_1}(a) = 0$

$\mathcal{A}_2 \models P(c)$

$D_{\mathcal{A}_2} = \{a\}$
 $c^{\mathcal{A}_2} = a$
 $P^{\mathcal{A}_2}(a) = 1$

$\mathcal{A}_3 \not\models P(c)$

$D_{\mathcal{A}_3} = \{b\}$
 $c^{\mathcal{A}_3} = b$
 $P^{\mathcal{A}_3}(b) = 0$

$\mathcal{A}_4 \models P(c)$

$D_{\mathcal{A}_4} = \{b\}$
 $c^{\mathcal{A}_4} = b$
 $P^{\mathcal{A}_4}(b) = 1$

$\mathcal{A}_5 \not\models P(c)$

$D_{\mathcal{A}_5} = \{a, b\}$
 $c^{\mathcal{A}_5} = b$
 $P^{\mathcal{A}_5}(b) = 0$
 $P^{\mathcal{A}_5}(a) = 0$
(or $P^{\mathcal{A}_5}(a) = 1$)

$\mathcal{A}_6 \models P(c)$

$D_{\mathcal{A}_6} = \{a, b\}$
 $c^{\mathcal{A}_6} = b$
 $P^{\mathcal{A}_6}(b) = 1$
 $P^{\mathcal{A}_6}(a) = 0$
(or $P^{\mathcal{A}_6}(a) = 1$)

Herbrand intuition

So many models! Even for a simple formula like $P(c)$ there are infinitely many structures and models

$\mathcal{A}_1 \not\models P(c)$

$D_{\mathcal{A}_1} = \{a\}$
 $c^{\mathcal{A}_1} = a$
 $P^{\mathcal{A}_1}(a) = 0$

$\mathcal{A}_2 \models P(c)$

$D_{\mathcal{A}_2} = \{a\}$
 $c^{\mathcal{A}_2} = a$
 $P^{\mathcal{A}_2}(a) = 1$

$\mathcal{A}_3 \not\models P(c)$

$D_{\mathcal{A}_3} = \{b\}$
 $c^{\mathcal{A}_3} = b$
 $P^{\mathcal{A}_3}(b) = 0$

$\mathcal{A}_4 \models P(c)$

$D_{\mathcal{A}_4} = \{b\}$
 $c^{\mathcal{A}_4} = b$
 $P^{\mathcal{A}_4}(b) = 1$

$\mathcal{A}_5 \not\models P(c)$

$D_{\mathcal{A}_5} = \{a, b\}$
 $c^{\mathcal{A}_5} = b$
 $P^{\mathcal{A}_5}(b) = 0$
 $P^{\mathcal{A}_5}(a) = 0$
(or $P^{\mathcal{A}_5}(a) = 1$)

$\mathcal{A}_6 \models P(c)$

$D_{\mathcal{A}_6} = \{a, b\}$
 $c^{\mathcal{A}_6} = b$
 $P^{\mathcal{A}_6}(b) = 1$
 $P^{\mathcal{A}_6}(a) = 0$
(or $P^{\mathcal{A}_6}(a) = 1$)

$\mathcal{A}_7 \not\models P(c)$

$D_{\mathcal{A}_7} = \{a, b\}$
 $c^{\mathcal{A}_7} = a$
 $P^{\mathcal{A}_7}(a) = 0$
 $P^{\mathcal{A}_7}(b) = 0$
(or $P^{\mathcal{A}_7}(b) = 1$)

Herbrand intuition

So many models! Even for a simple formula like $P(c)$ there are infinitely many structures and models

$\mathcal{A}_1 \not\models P(c)$

$D_{\mathcal{A}_1} = \{a\}$
 $c^{\mathcal{A}_1} = a$
 $P^{\mathcal{A}_1}(a) = 0$

$\mathcal{A}_2 \models P(c)$

$D_{\mathcal{A}_2} = \{a\}$
 $c^{\mathcal{A}_2} = a$
 $P^{\mathcal{A}_2}(a) = 1$

$\mathcal{A}_3 \not\models P(c)$

$D_{\mathcal{A}_3} = \{b\}$
 $c^{\mathcal{A}_3} = b$
 $P^{\mathcal{A}_3}(b) = 0$

$\mathcal{A}_4 \models P(c)$

$D_{\mathcal{A}_4} = \{b\}$
 $c^{\mathcal{A}_4} = b$
 $P^{\mathcal{A}_4}(b) = 1$

$\mathcal{A}_5 \not\models P(c)$

$D_{\mathcal{A}_5} = \{a, b\}$
 $c^{\mathcal{A}_5} = b$
 $P^{\mathcal{A}_5}(b) = 0$
 $P^{\mathcal{A}_5}(a) = 0$
(or $P^{\mathcal{A}_5}(a) = 1$)

$\mathcal{A}_6 \models P(c)$

$D_{\mathcal{A}_6} = \{a, b\}$
 $c^{\mathcal{A}_6} = b$
 $P^{\mathcal{A}_6}(b) = 1$
 $P^{\mathcal{A}_6}(a) = 0$
(or $P^{\mathcal{A}_6}(a) = 1$)

$\mathcal{A}_7 \not\models P(c)$

$D_{\mathcal{A}_7} = \{a, b\}$
 $c^{\mathcal{A}_7} = a$
 $P^{\mathcal{A}_7}(a) = 0$
 $P^{\mathcal{A}_7}(b) = 0$
(or $P^{\mathcal{A}_7}(b) = 1$)

$\mathcal{A}_8 \models P(c)$

$D_{\mathcal{A}_8} = \{a, b\}$
 $c^{\mathcal{A}_8} = a$
 $P^{\mathcal{A}_8}(a) = 1$
 $P^{\mathcal{A}_8}(b) = 0$
(or $P^{\mathcal{A}_8}(b) = 1$)

Herbrand intuition

So many models! Even for a simple formula like $P(c)$ there are infinitely many structures and models

$\mathcal{A}_1 \not\models P(c)$

$D_{\mathcal{A}_1} = \{a\}$
 $c^{\mathcal{A}_1} = a$
 $P^{\mathcal{A}_1}(a) = 0$

$\mathcal{A}_2 \models P(c)$

$D_{\mathcal{A}_2} = \{a\}$
 $c^{\mathcal{A}_2} = a$
 $P^{\mathcal{A}_2}(a) = 1$

$\mathcal{A}_3 \not\models P(c)$

$D_{\mathcal{A}_3} = \{b\}$
 $c^{\mathcal{A}_3} = b$
 $P^{\mathcal{A}_3}(b) = 0$

$\mathcal{A}_4 \models P(c)$

$D_{\mathcal{A}_4} = \{b\}$
 $c^{\mathcal{A}_4} = b$
 $P^{\mathcal{A}_4}(b) = 1$

$\mathcal{A}_5 \not\models P(c)$

$D_{\mathcal{A}_5} = \{a, b\}$
 $c^{\mathcal{A}_5} = b$
 $P^{\mathcal{A}_5}(b) = 0$
 $P^{\mathcal{A}_5}(a) = 0$
(or $P^{\mathcal{A}_5}(a) = 1$)

$\mathcal{A}_6 \models P(c)$

$D_{\mathcal{A}_6} = \{a, b\}$
 $c^{\mathcal{A}_6} = b$
 $P^{\mathcal{A}_6}(b) = 1$
 $P^{\mathcal{A}_6}(a) = 0$
(or $P^{\mathcal{A}_6}(a) = 1$)

$\mathcal{A}_7 \not\models P(c)$

$D_{\mathcal{A}_7} = \{a, b\}$
 $c^{\mathcal{A}_7} = a$
 $P^{\mathcal{A}_7}(a) = 0$
 $P^{\mathcal{A}_7}(b) = 0$
(or $P^{\mathcal{A}_7}(b) = 1$)

$\mathcal{A}_8 \models P(c)$

$D_{\mathcal{A}_8} = \{a, b\}$
 $c^{\mathcal{A}_8} = a$
 $P^{\mathcal{A}_8}(a) = 1$
 $P^{\mathcal{A}_8}(b) = 0$
(or $P^{\mathcal{A}_8}(b) = 1$)

- All structures are quite similar!

Herbrand intuition

So many models! Even for a simple formula like $P(c)$ there are infinitely many structures and models

$\mathcal{A}_1 \not\models P(c)$

$D_{\mathcal{A}_1} = \{a\}$
 $c^{\mathcal{A}_1} = a$
 $P^{\mathcal{A}_1}(a) = 0$

$\mathcal{A}_2 \models P(c)$

$D_{\mathcal{A}_2} = \{a\}$
 $c^{\mathcal{A}_2} = a$
 $P^{\mathcal{A}_2}(a) = 1$

$\mathcal{A}_3 \not\models P(c)$

$D_{\mathcal{A}_3} = \{b\}$
 $c^{\mathcal{A}_3} = b$
 $P^{\mathcal{A}_3}(b) = 0$

$\mathcal{A}_4 \models P(c)$

$D_{\mathcal{A}_4} = \{b\}$
 $c^{\mathcal{A}_4} = b$
 $P^{\mathcal{A}_4}(b) = 1$

$\mathcal{A}_5 \not\models P(c)$

$D_{\mathcal{A}_5} = \{a, b\}$
 $c^{\mathcal{A}_5} = b$
 $P^{\mathcal{A}_5}(b) = 0$
 $P^{\mathcal{A}_5}(a) = 0$
(or $P^{\mathcal{A}_5}(a) = 1$)

$\mathcal{A}_6 \models P(c)$

$D_{\mathcal{A}_6} = \{a, b\}$
 $c^{\mathcal{A}_6} = b$
 $P^{\mathcal{A}_6}(b) = 1$
 $P^{\mathcal{A}_6}(a) = 0$
(or $P^{\mathcal{A}_6}(a) = 1$)

$\mathcal{A}_7 \not\models P(c)$

$D_{\mathcal{A}_7} = \{a, b\}$
 $c^{\mathcal{A}_7} = a$
 $P^{\mathcal{A}_7}(a) = 0$
 $P^{\mathcal{A}_7}(b) = 0$
(or $P^{\mathcal{A}_7}(b) = 1$)

$\mathcal{A}_8 \models P(c)$

$D_{\mathcal{A}_8} = \{a, b\}$
 $c^{\mathcal{A}_8} = a$
 $P^{\mathcal{A}_8}(a) = 1$
 $P^{\mathcal{A}_8}(b) = 0$
(or $P^{\mathcal{A}_8}(b) = 1$)

- All structures are quite similar!
- Changing domains does not seem to change much

Herbrand intuition

So many models! Even for a simple formula like $P(c)$ there are infinitely many structures and models

$\mathcal{A}_1 \not\models P(c)$

$D_{\mathcal{A}_1} = \{a\}$
 $c^{\mathcal{A}_1} = a$
 $P^{\mathcal{A}_1}(a) = 0$

$\mathcal{A}_2 \models P(c)$

$D_{\mathcal{A}_2} = \{a\}$
 $c^{\mathcal{A}_2} = a$
 $P^{\mathcal{A}_2}(a) = 1$

$\mathcal{A}_3 \not\models P(c)$

$D_{\mathcal{A}_3} = \{b\}$
 $c^{\mathcal{A}_3} = b$
 $P^{\mathcal{A}_3}(b) = 0$

$\mathcal{A}_4 \models P(c)$

$D_{\mathcal{A}_4} = \{b\}$
 $c^{\mathcal{A}_4} = b$
 $P^{\mathcal{A}_4}(b) = 1$

$\mathcal{A}_5 \not\models P(c)$

$D_{\mathcal{A}_5} = \{a, b\}$
 $c^{\mathcal{A}_5} = b$
 $P^{\mathcal{A}_5}(b) = 0$
 $P^{\mathcal{A}_5}(a) = 0$
(or $P^{\mathcal{A}_5}(a) = 1$)

$\mathcal{A}_6 \models P(c)$

$D_{\mathcal{A}_6} = \{a, b\}$
 $c^{\mathcal{A}_6} = b$
 $P^{\mathcal{A}_6}(b) = 1$
 $P^{\mathcal{A}_6}(a) = 0$
(or $P^{\mathcal{A}_6}(a) = 1$)

$\mathcal{A}_7 \not\models P(c)$

$D_{\mathcal{A}_7} = \{a, b\}$
 $c^{\mathcal{A}_7} = a$
 $P^{\mathcal{A}_7}(a) = 0$
 $P^{\mathcal{A}_7}(b) = 0$
(or $P^{\mathcal{A}_7}(b) = 1$)

$\mathcal{A}_8 \models P(c)$

$D_{\mathcal{A}_8} = \{a, b\}$
 $c^{\mathcal{A}_8} = a$
 $P^{\mathcal{A}_8}(a) = 1$
 $P^{\mathcal{A}_8}(b) = 0$
(or $P^{\mathcal{A}_8}(b) = 1$)

- All structures are quite similar!
- Changing domains does not seem to change much
- The interpretation of predicates appears crucial

Herbrand intuition

So many models! Even for a simple formula like $P(c)$ there are infinitely many structures and models

$\mathcal{A}_1 \not\models P(c)$

$D_{\mathcal{A}_1} = \{a\}$
 $c^{\mathcal{A}_1} = a$
 $P^{\mathcal{A}_1}(a) = 0$

$\mathcal{A}_2 \models P(c)$

$D_{\mathcal{A}_2} = \{a\}$
 $c^{\mathcal{A}_2} = a$
 $P^{\mathcal{A}_2}(a) = 1$

$\mathcal{A}_3 \not\models P(c)$

$D_{\mathcal{A}_3} = \{b\}$
 $c^{\mathcal{A}_3} = b$
 $P^{\mathcal{A}_3}(b) = 0$

$\mathcal{A}_4 \models P(c)$

$D_{\mathcal{A}_4} = \{b\}$
 $c^{\mathcal{A}_4} = b$
 $P^{\mathcal{A}_4}(b) = 1$

$\mathcal{A}_5 \not\models P(c)$

$D_{\mathcal{A}_5} = \{a, b\}$
 $c^{\mathcal{A}_5} = b$
 $P^{\mathcal{A}_5}(b) = 0$
 $P^{\mathcal{A}_5}(a) = 0$
(or $P^{\mathcal{A}_5}(a) = 1$)

$\mathcal{A}_6 \models P(c)$

$D_{\mathcal{A}_6} = \{a, b\}$
 $c^{\mathcal{A}_6} = b$
 $P^{\mathcal{A}_6}(b) = 1$
 $P^{\mathcal{A}_6}(a) = 0$
(or $P^{\mathcal{A}_6}(a) = 1$)

$\mathcal{A}_7 \not\models P(c)$

$D_{\mathcal{A}_7} = \{a, b\}$
 $c^{\mathcal{A}_7} = a$
 $P^{\mathcal{A}_7}(a) = 0$
 $P^{\mathcal{A}_7}(b) = 0$
(or $P^{\mathcal{A}_7}(b) = 1$)

$\mathcal{A}_8 \models P(c)$

$D_{\mathcal{A}_8} = \{a, b\}$
 $c^{\mathcal{A}_8} = a$
 $P^{\mathcal{A}_8}(a) = 1$
 $P^{\mathcal{A}_8}(b) = 0$
(or $P^{\mathcal{A}_8}(b) = 1$)

- All structures are quite similar!
- Changing domains does not seem to change much
- The interpretation of predicates appears crucial
- The interpretation of functions appears to be **isomorphic** for different domains

Cardinality of domain

- Changing domains does not seem to change much

Cardinality of domain

- Changing domains does not seem to change much
- But...

$$P(c) \wedge \neg P(d)$$

Cardinality of domain

- Changing domains does not seem to change much
- But...

$$P(c) \wedge \neg P(d)$$

$$\mathcal{A} \models P(c) \wedge \neg P(d)$$

$$D_{\mathcal{A}} = \{a, b\}$$

$$c^{\mathcal{A}} = a$$

$$d^{\mathcal{A}} = b$$

$$P^{\mathcal{A}}(a) = 1$$

$$P^{\mathcal{A}}(b) = 0$$

Cardinality of domain

- Changing domains does not seem to change much
- But...

$$P(c) \wedge \neg P(d)$$

$$\mathcal{A} \models P(c) \wedge \neg P(d)$$

$$D_{\mathcal{A}} = \{a, b\}$$

$$c^{\mathcal{A}} = a$$

$$d^{\mathcal{A}} = b$$

$$P^{\mathcal{A}}(a) = 1$$

$$P^{\mathcal{A}}(b) = 0$$

- Is there a model whose domain has cardinality 1?

Cardinality of domain

- Changing domains does not seem to change much
- But...

$$P(c) \wedge \neg P(d)$$

$$\mathcal{A} \models P(c) \wedge \neg P(d)$$

$$D_{\mathcal{A}} = \{a, b\}$$

$$c^{\mathcal{A}} = a$$

$$d^{\mathcal{A}} = b$$

$$P^{\mathcal{A}}(a) = 1$$

$$P^{\mathcal{A}}(b) = 0$$

- Is there a model whose domain has cardinality 1?
- No, there isn't!

Cardinality of domain

- Changing domains does not seem to change much
- But...

$$P(c) \wedge \neg P(d)$$

$$\mathcal{A} \models P(c) \wedge \neg P(d)$$

$$D_{\mathcal{A}} = \{a, b\}$$

$$c^{\mathcal{A}} = a$$

$$d^{\mathcal{A}} = b$$

$$P^{\mathcal{A}}(a) = 1$$

$$P^{\mathcal{A}}(b) = 0$$

- Is there a model whose domain has cardinality 1?
- No, there isn't!
- Cardinality of the domain is important

- 1 Normal forms
 - Prenex Normal Form
 - Skolemization
- 2 Herbrand theory
 - Intuition
 - **Main statement**
- 3 Exercises

Herbrand interpretation — \mathcal{H}

- Use the set of **ground terms** of the formula as domain!
 - This domain is called **Herbrand universe**
 - **Note:** If the language contains no constant, add an arbitrary constant

Herbrand interpretation — \mathcal{H}

- Use the set of **ground terms** of the formula as domain!
 - This domain is called **Herbrand universe**
 - **Note:** If the language contains no constant, add an arbitrary constant
- Interpret function symbols as **themselves**
 - $c^{\mathcal{H}} = c$ for constants
 - $f^{\mathcal{H}}(t_1, \dots, t_n) = f(t_1^{\mathcal{H}}, \dots, t_n^{\mathcal{H}})$

Herbrand interpretation — \mathcal{H}

- Use the set of **ground terms** of the formula as domain!
 - This domain is called **Herbrand universe**
 - **Note:** If the language contains no constant, add an arbitrary constant
- Interpret function symbols as **themselves**
 - $c^{\mathcal{H}} = c$ for constants
 - $f^{\mathcal{H}}(t_1, \dots, t_n) = f(t_1^{\mathcal{H}}, \dots, t_n^{\mathcal{H}})$
- **Note:** Interpretations of predicates is not fixed

Herbrand interpretation — \mathcal{H}

- Use the set of **ground terms** of the formula as domain!
 - This domain is called **Herbrand universe**
 - **Note:** If the language contains no constant, add an arbitrary constant
- Interpret function symbols as **themselves**
 - $c^{\mathcal{H}} = c$ for constants
 - $f^{\mathcal{H}}(t_1, \dots, t_n) = f(t_1^{\mathcal{H}}, \dots, t_n^{\mathcal{H}})$
- **Note:** Interpretations of predicates is not fixed
- **Herbrand base:** $\{P(t_1, \dots, t_n) \mid P \text{ is a predicate of arity } n, t_1, \dots, t_n \text{ are ground terms}\}$

Herbrand interpretation — \mathcal{H}

- Use the set of **ground terms** of the formula as domain!
 - This domain is called **Herbrand universe**
 - **Note:** If the language contains no constant, add an arbitrary constant
- Interpret function symbols as **themselves**
 - $c^{\mathcal{H}} = c$ for constants
 - $f^{\mathcal{H}}(t_1, \dots, t_n) = f(t_1^{\mathcal{H}}, \dots, t_n^{\mathcal{H}})$
- **Note:** Interpretations of predicates is not fixed
- **Herbrand base:** $\{P(t_1, \dots, t_n) \mid P \text{ is a predicate of arity } n, t_1, \dots, t_n \text{ are ground terms}\}$
- Each predicate is a subset of the Herbrand base

Herbrand interpretation — \mathcal{H}

- Use the set of **ground terms** of the formula as domain!
 - This domain is called **Herbrand universe**
 - **Note:** If the language contains no constant, add an arbitrary constant
- Interpret function symbols as **themselves**
 - $c^{\mathcal{H}} = c$ for constants
 - $f^{\mathcal{H}}(t_1, \dots, t_n) = f(t_1^{\mathcal{H}}, \dots, t_n^{\mathcal{H}})$
- **Note:** Interpretations of predicates is not fixed
- **Herbrand base:** $\{P(t_1, \dots, t_n) \mid P \text{ is a predicate of arity } n, t_1, \dots, t_n \text{ are ground terms}\}$
- Each predicate is a subset of the Herbrand base

An Herbrand model of a set Γ of wffs is an Herbrand interpretation satisfying Γ

Example

- 1 $N(o)$
 - 2 $\forall x (N(x) \rightarrow N(s(x)))$
 - 3 $\forall x \forall y (\neg E(x, y) \rightarrow \neg E(s(x), s(y)))$
 - 4 $\forall x \neg E(s(x), o)$
-
- $D_{\mathcal{H}} = \{o, s(o), s(s(o)), s(s(s(o))), \dots\}$
 - $o^{\mathcal{A}} = o$
 - $s^{\mathcal{A}}(o) = s(o); s^{\mathcal{A}}(s(o)) = s(s(o));$
 $s^{\mathcal{A}}(s(s(o))) = s(s(s(o))); \dots$
 - $N^{\mathcal{A}}(o) = 1; N^{\mathcal{A}}(s(o)) = 1; N^{\mathcal{A}}(s(s(o))) = 1; \dots$
 - $E^{\mathcal{A}}(o, o) = 1; E^{\mathcal{A}}(o, s(o)) = 0; E^{\mathcal{A}}(s(o), o) = 0;$
 $E^{\mathcal{A}}(s(o), s(o)) = 1; E^{\mathcal{A}}(s(o), s(s(o))) = 0; \dots$

Theorem

*Let Γ be a set of closed wffs in Skolem normal form.
 Γ is satisfiable if and only if Γ has a Herbrand model.*

Sketch. (\Leftarrow) Immediate.

(\Rightarrow) Use structural induction. For the universal quantifier use the following lemma:

Translation Lemma

For any wff ϕ we have

$$\nu^{(\mathcal{A}, \xi^{\mathcal{A}})}(\phi[t/x]) = \nu^{(\mathcal{A}, \xi^{\mathcal{A}}[\nu^{(\mathcal{A}, \xi^{\mathcal{A}})}(t)/x])}(\phi)$$

Read as follows: The evaluation of $\phi[t/x]$ wrt $(\mathcal{A}, \xi^{\mathcal{A}})$ is equal to the evaluation of ϕ wrt $(\mathcal{A}, \xi^{\mathcal{A}}[\nu^{(\mathcal{A}, \xi^{\mathcal{A}})}(t)/x])$

- We cannot check all Herbrand interpretations!

- We cannot check all Herbrand interpretations!
- We were actually interested in modeling unsatisfiability

- We cannot check all Herbrand interpretations!
- We were actually interested in modeling unsatisfiability

Definition

Let $\phi = \forall x_1 \cdots \forall x_n \psi$ be a formula in Skolem normal form. The Herbrand expansion of ϕ , denoted $\varepsilon(\phi)$, is the following set of ground formulas:

$$\varepsilon(\phi) = \{\psi[t_1/x_1, \dots, t_n/x_n] \mid t_1, \dots, t_n \in D_{\mathcal{H}}\}$$

- We cannot check all Herbrand interpretations!
- We were actually interested in modeling unsatisfiability

Definition

Let $\phi = \forall x_1 \cdots \forall x_n \psi$ be a formula in Skolem normal form. The Herbrand expansion of ϕ , denoted $\varepsilon(\phi)$, is the following set of ground formulas:

$$\varepsilon(\phi) = \{\psi[t_1/x_1, \dots, t_n/x_n] \mid t_1, \dots, t_n \in D_{\mathcal{H}}\}$$

- $\varepsilon(\phi)$ is a set of propositional wffs

- We cannot check all Herbrand interpretations!
- We were actually interested in modeling unsatisfiability

Definition

Let $\phi = \forall x_1 \cdots \forall x_n \psi$ be a formula in Skolem normal form. The Herbrand expansion of ϕ , denoted $\varepsilon(\phi)$, is the following set of ground formulas:

$$\varepsilon(\phi) = \{\psi[t_1/x_1, \dots, t_n/x_n] \mid t_1, \dots, t_n \in D_{\mathcal{H}}\}$$

- $\varepsilon(\phi)$ is a set of propositional wffs
- ϕ is satisfiable if and only if $\varepsilon(\phi)$ is satisfiable

- We cannot check all Herbrand interpretations!
- We were actually interested in modeling unsatisfiability

Definition

Let $\phi = \forall x_1 \cdots \forall x_n \psi$ be a formula in Skolem normal form. The Herbrand expansion of ϕ , denoted $\varepsilon(\phi)$, is the following set of ground formulas:

$$\varepsilon(\phi) = \{\psi[t_1/x_1, \dots, t_n/x_n] \mid t_1, \dots, t_n \in D_{\mathcal{H}}\}$$

- $\varepsilon(\phi)$ is a set of propositional wffs
- ϕ is satisfiable if and only if $\varepsilon(\phi)$ is satisfiable
- By the Compactness Theorem, $\varepsilon(\phi)$ is unsatisfiable if and only if there is a finite subset of $\varepsilon(\phi)$ that is unsatisfiable

- Let ϕ be a formula in Skolem normal form
- Fix an enumeration for the elements of $\varepsilon(\phi)$

Algorithm

- 1 $n := 0$
- 2 $n := n + 1$
- 3 If $\phi_1 \wedge \dots \wedge \phi_n$ is unsatisfiable then output ϕ is unsatisfiable
- 4 Go to 2

- Let ϕ be a formula in Skolem normal form
- Fix an enumeration for the elements of $\varepsilon(\phi)$

Algorithm

- 1 $n := 0$
- 2 $n := n + 1$
- 3 If $\phi_1 \wedge \dots \wedge \phi_n$ is unsatisfiable then output ϕ is unsatisfiable
- 4 Go to 2

- How to check unsatisfiability of a set Γ of wffs?


- 1 Normal forms
 - Prenex Normal Form
 - Skolemization
- 2 Herbrand theory
 - Intuition
 - Main statement
- 3 Exercises

- 1 Transform the formula

$$\neg \exists u ((\forall x P(x, u)) \rightarrow (\forall y \exists z (\forall v Q(u, v, y, z) \wedge \forall w Q(w, u, z, y))))$$

into Skolem Normal Form

- 2 Solve Exercise 3.8, 3.9, 3.10 and 3.15 in the booklet of Ghidini and Serafini (not using the solution!)
- 3 Exercises from 3.16 to the end of the chapter ask to model a scenario. Have a look at them!



END OF THE
LECTURE